

Literature Citing Actim[®] Partus



The list below includes examples of literature that reference Actim[®] Partus, and serves as an illustrative sample of its presence in published research.

Published Studies Referencing Actim[®] Partus

Rui-Hong Lan et al. Significance of highly phosphorylated insulin-like growth factor binding protein-1 and cervical length for prediction of preterm delivery in twin pregnancies. *World J Clin Cases* (2021) **9**:4553-4558

Chen MX, Dansereau J, Hoag GN. Comparison of Fetal Fibronectin and Phosphorylated Insulin-Like Growth Factor Binding Protein-1 Testing to Predict Preterm Delivery in Symptomatic Women: A 10-Year Retrospective Study. *J Obstet Gynaecol Can.* (2020) **42**:971-976

Harley-Campbell J et al. Three biomarker tests to help diagnose preterm labour: a systematic review and economic evaluation. *Health Technol Assess.* (2019) **23**:1-226.

Vallikkannu N et al. Insulin-like growth factor binding protein 1, Bishop score, and sonographic cervical length: tolerability and prediction of vaginal birth and vaginal birth within 24 hours following labour induction in nulliparous women. *BJOG* (2017) **124**:1274-1283.

Bruijn MM et al. Comparison of the Actim Partus test and the fetal fibronectin test in the prediction of spontaneous preterm birth in symptomatic women undergoing cervical length measurement. *Eur J Obstet Gynecol Reprod Biol.* (2016) **206**:220-224.

Đogić et al. IGFBP-1 marker of cervical ripening and predictor of preterm birth. *Medicinski Glasnik* (2016) **13**.

Tripathi R et al. Comparison of rapid bedside tests for phosphorylated insulin-like growth factor-binding protein 1 and fetal fibronectin to predict preterm birth. *Int J Gynaecol Obstet.* (2016) **135**:47-50. Epub 2016 Jun 18.

Cooper S et al. Diagnostic accuracy of rapid pHIGFBP-1 assay for predicting preterm labor in symptomatic patients. *J Perinatol.* (2012) **32**:460-465. Epub 2011 Oct 13.

Danti L et al. The combination of short cervical length and pHIGFBP-1 in the prediction of preterm delivery in symptomatic women. *J Matern Fetal Neonatal Med.* (2011) **24**:1262-1266.

Dögl M, Skogvoll E, Heimstad R. Cervical insulin-like growth factor binding protein-1 (IGFBP-1) to predict spontaneous onset of labor and induction to delivery interval in post-term pregnancy. *Acta Obstet Gynecol Scand.* (2011) **90**:57-62.

Rahkonen L. Prediction of pre-term delivery with phosphorylated insulin-like growth factor-binding protein-1. *European Obstetrics & Gynecology* (2011) **6**:3-7.

Riboni F et al. Biochemical markers predicting pre-term delivery in symptomatic patients: phosphorylated insulin-like growth factor binding protein-1 and fetal fibronectin. *Arch Gynecol Obstet.* (2011) **284**:1325-9.

Riboni F et al. Combination of biochemical markers in predicting pre-term delivery. *Arch Gynecol Obstet.* (2012) **285**:61-66.

Adeyemi O, Osoba L. The role of phosphorylated insulin-like growth factor binding protein-1 in predicting pre-term labour in twin pregnancies. *Journal of Obstetrics and Gynaecology* (2010) **30**:571-573.

Brik Spinelli M et al. Phosphorylated insulin-like growth factor binding protein-1 and cervical measurement in women with threatening preterm birth. *Acta Obstet Gynecol Scand* (2010) **89**:268-74.

Nor Azlin MI et al. Role of pHIGFBP-1 and ultrasound cervical length in predicting pre-term labour. *Journal of Obstetrics and Gynaecology* (2010) **30**:456-460.

Tanir HM, Sener T, Yildiz Z. Cervical phosphorylated insulin-like growth factor binding protein-1 for the prediction of preterm delivery in symptomatic cases with intact membranes. *J Obstet Gynaecol Res* (2009) **1**:66-72.

Altinkaya O et al. Cervical phosphorylated insulin-like growth factor binding protein-1 in prediction of preterm delivery. *Arch Gynecol Obstet* (2009) **279**:279-283. Epub 2008 Jun 13.

Balic D, Latifagic A, Hudic I. Insulin-like growth factor-binding protein-1 (IGFBP-1) in cervical secretions as a predictor of preterm delivery. *J Matern Fetal Neonatal Med* (2008) **21**:297-300.

Bittar R et al. Predicting preterm delivery in asymptomatic patients with prior preterm delivery by measurement of cervical length and phosphorylated insulin-like growth factor-binding protein-1. *Ultrasound Obstet Gynecol* (2007) **29**:562-567.

Eroglu D et al. Prediction of preterm delivery among women with threatened preterm labor. *Gynecol Obstet Invest* (2007) **64**:109-116.

Paternoster DM et al. Cervical pHIGFBP-1 in the evaluation of the risk of preterm delivery. *Acta Obstet Gynecol Scand* (2007) **86**:151-155.

Ting HS et al. Comparison of bedside test kits for prediction of preterm delivery: phosphorylated insulin-like growth factor binding protein-1 (pIGFBP-1) test and fetal fibronectin test. *Ann Acad Med Singapore* (2007) **36**:399-402.

Elizur SE et al. Insulin-like growth factor binding protein-1 detection in preterm labor: evaluation of a bedside test. *Am J Perinatol* (2005) **22**:305-309.

Akercan F et al. Value of cervical phosphorylated insulinlike growth factor binding protein-1 in the prediction of preterm labor. *J Reprod Med* (2004) **49**:368-372.

Kwek K et al. Evaluation of a bedside test for phosphorylated insulin-like growth factor binding protein-1 in preterm labour. *Ann Acad Med Singapore* (2004) **33**:780-783.

Park O-R et al. Usefulness of phosphorylated insulin-like growth factor binding protein-1 for prediction of preterm delivery. *Korean J Obstet Gynecol* (2003) **46**:1378-1383. (In Korean, abstract in English).

Lembet A et al. New rapid bed-side test to predict preterm delivery: phosphorylated insulin-like growth factor binding protein-1 in cervical secretions. *Acta Obstet Gynecol Scand* (2002) **81**:706-712.

Kekki M et al. Insulin-like growth factor-binding protein-1 in cervical secretion as a predictor of preterm delivery. *Acta Obstet Gynecol Scand* (2001) **80**:546-551.

Shine BK et al. Insulin-like growth factor-binding protein-1 in cervical secretion as a predictor of preterm delivery. *Korean J Obstet Gynecol* (2001) **44**:2250-2256. (In Korean, abstract in English).

Kekki M et al. Insulin-like growth factor binding protein-1 in cervix as a marker of infectious complications in pregnant women with bacterial vaginosis. *Lancet* (1999) **353**:1494.

Nuutila M et al. Phosphorylated isoforms of insulin-like growth factor binding protein-1 in the cervix as a predictor of cervical ripeness. *Obstet Gynecol* (1999) **94**:243-249.

Abstracts & Posters on Actim[®] Partus

Sanchez Martinez M et al. Comparison of ecographic cervical length at two different cut-off points and two biochemical markers as predictors of spontaneous preterm delivery in women admitted because of preterm labor. *Ultrasound Obstet Gynecol* (2006) **28**:583-584. Poster abstract in 16th World Congress on Ultrasound in Obstetrics and Gynecology, Sept. 3-7, 2006, London, UK.

Turnell R et al. A direct comparison of fetal fibronectin [FFN@] and pIGFBP-1 [actim[™] Partus] in the diagnosis of preterm labour [PTL]. Abstract in JOGC (2005), **17**:O-OBS-014.

Bittar R et al. Cervical insulin like growth factor binding protein-1 (pHIGFBP-1) in patients at increased risk for preterm delivery: preliminary results. Poster presented in 5th World Congress of Perinatal Medicine, Sept. 23-27, 2001, Barcelona, Spain.

Articles on IGFBP-1

Rutanen E-M. Insulin-like growth factors in obstetrics. *Opin Obstet Gynecol* (2000) **12**:163-168.